

ON QUANTUM DE RHAM COHOMOLOGY THEORY

HUI-DONG CAO & JIAN ZHOU

ABSTRACT. We define quantum exterior product $\wedge_h$ and quantum exterior differential d_h on Poisson manifolds (of which symplectic manifolds are an important class of examples). Quantum de Rham cohomology, which is a deformation quantization of de Rham cohomology, is defined as the cohomology of d_h . We also define quantum Dolbeault cohomology. A version of quantum integral on symplectic manifolds is considered and the corresponding quantum Stokes theorem is proved. We also derive quantum hard Lefschetz theorem. By replacing d by d_h and $\wedge$ by $\wedge_h$ in the usual definitions, we define many quantum analogues of important objects in differential geometry, e.g. quantum curvature. The quantum characteristic classes are then studied along the lines of classical Chern-Weil theory. Quantum equivariant de Rham cohomology is defined in the similar fashion.

Recently, quantum cohomology has attracted a lot of attentions. See Tian [8] for a survey on this topic, and the introduction of Li-Tian [7] for more recent development. In this note we announce a new construction of a deformation of the ring structure on the de Rham cohomology of a closed symplectic manifold. Our construction is of algebraic nature, following the definition of the de Rham cohomology. In fact, on any Poisson manifold, we will define a quantum exterior product $\wedge_h$ of exterior forms, and quantum exterior differential d_h , such that $d_h^2 = 0$, and d_h is a derivation for $\wedge_h$. We define *quantum de Rham cohomology* as the cohomology of d_h . By replacing d by d_h and $\wedge$ by $\wedge_h$ in the usual definitions, we define many quantum analogues of important objects, such as quantum curvature and quantum characteristic classes, in differential geometry.

Our construction has the following features which are not shared by the quantum cohomology:

1. The proof of the associativity of the quantum exterior product is elementary.
2. It is routine to define quantum Dolbeault cohomology.
3. It is routine to define quantum characteristic class.
4. It is routine to define quantum equivariant de Rham cohomology.
5. The computations for homogeneous examples are elementary.

The definition of our quantum exterior product is motivated by Moyal-Weyl multiplication and Clifford multiplication. For any finite dimensional vector space V , let $\{e_1, \dots, e_m\}$ be a basis of V and $\{e^1, \dots, e^m\}$ be the dual basis. Assume that $w = w^{ij}e_i \otimes e_j \in V \otimes V$. Then w defines a multiplication $\wedge_w$ on $\Lambda(V^*)$, and a multiplication $*_w$ on $S(V^*)$, such that $e^i \wedge_w e^j = e^i \wedge e^j + w^{ij}$, $e^i *_w e^j = e^i \odot e^j + w^{ij}$. If $w \in S^2(V)$, then $\wedge_w$ is the Clifford multiplication. If $w \in \Lambda^2(V)$ is nondegenerate, $*_w$ is the Moyal-Weyl multiplication. If $w \in \Lambda^2(V)$, then $\wedge_w$ is what we call the *quantum exterior product* (or a quantum Clifford multiplication). It is elementary

¹1991 *Mathematics Subject Classification*:

²Authors' research was supported in part by NSF

to show that this multiplication is associative. We use it to obtain a quantum calculus on any Poisson manifold. We will present the main results below, and the proofs will appear elsewhere.

1. Quantum exterior algebra. Let V be a finite dimensional vector space over a field $\mathbf{k}$ of characteristic zero, and $\Lambda(V^*)$ be the exterior algebra generated by the dual vector space V^* . For any $v \in V$ and $\alpha \in \Lambda^k(V^*)$, denote

$$\begin{aligned} (v \vdash \alpha)(v_1, \dots, v_{k-1}) &= \alpha(v, v_1, \dots, v_{k-1}), \\ (\alpha \dashv v)(v_1, \dots, v_{k-1}) &= \alpha(v_1, \dots, v_{k-1}, v), \end{aligned}$$

for $v_1, \dots, v_{k-1} \in V$. Let $\Lambda_h(V^*) = \Lambda(V^*)[h] = \Lambda(V^*) \otimes_{\mathbf{k}} \mathbf{k}[h]$. For any $w \in \Lambda^2(V)$, with $w = w^{ij} e_i \wedge e_j$ with respect to a basis $\{e_1, \dots, e_m\}$ of V , we define the *quantum exterior product* $\wedge_{h,w} : \Lambda(V^*) \otimes \Lambda(V^*) \rightarrow \Lambda(V^*)[h]$ by

$$\begin{aligned} &\alpha \wedge_{h,w} \beta \\ &= \sum_{n \geq 0} \frac{h^n}{n!} w^{i_1 j_1} \dots w^{i_n j_n} (\alpha \dashv e_{i_1} \dashv \dots \dashv e_{i_n}) \wedge (e_{j_n} \vdash \dots \vdash e_{j_1} \vdash \beta), \end{aligned}$$

for $\alpha, \beta \in \Lambda(V^*)$. This definition is evidently independent of the choice of the basis. We extend $\wedge_{h,w}$ as $\mathbf{k}[h]$ -module map to $\Lambda_h(V^*) \otimes_{\mathbf{k}[h]} \Lambda_h(V^*)$. When there is no confusion about w , we will simply write $\alpha \wedge_h \beta$ for $\alpha \wedge_{h,w} \beta$. We are interested in $\alpha \wedge_w \beta$, which is just $\alpha \wedge_{1,w} \beta$. We assign h the degree 2. Then $\Lambda_h(V^*)$ has a natural $\mathbb{Z}$ -grading. Denote by $\Lambda_h^{[n]}(V^*)$ the subspace of homogeneous elements of degree n , then it is clear that

$$\Lambda_h^{[m]}(V^*) \wedge_h \Lambda_h^{[n]}(V^*) \subset \Lambda_h^{[m+n]}(V^*).$$

Theorem 1. (Cao-Zhou [3]) *The quantum exterior product satisfies the following properties*

- (1) *Supercommutativity* $\alpha \wedge_h \beta = (-1)^{|\alpha||\beta|} \beta \wedge_h \alpha,$
- (2) *Associativity* $(\alpha \wedge_h \beta) \wedge_h \gamma = \alpha \wedge_h (\beta \wedge_h \gamma),$

for all $\alpha, \beta, \gamma \in \Lambda_h(V^*)$. Therefore, $(\Lambda_h(V^*), \wedge_h)$ is a deformation quantization of the exterior algebra $(\Lambda(V^*), \wedge)$.

The proof of (1) is trivial. The proof of (2) is of elementary nature but non-trivial. It is proved first by brute force in the case of $\deg(\alpha) = 1$, then by induction on $\deg(\alpha)$ (see [3] for details).

We can also extend $\wedge_h$ to $\Lambda_{h,h^{-1}}(V^*) = \Lambda(V^*)[h, h^{-1}] = \Lambda(V^*) \otimes_{\mathbf{k}} \mathbf{k}[h, h^{-1}]$.

An algebra A with unit $e \in A$ over a field $\mathbf{k}$ is called a **$\mathbf{k}$ -Fröbenius algebra** if there is a nondegenerate symmetric $\mathbf{k}$ -bilinear function $\langle \cdot, \cdot \rangle : A \times A \rightarrow \mathbf{k}$, such that

$$\langle \alpha \beta, \gamma \rangle = \langle \alpha, \beta \gamma \rangle,$$

for all $\alpha, \beta, \gamma \in A$. There is a simple way to construct a structure of Fröbenius algebra on any $\mathbf{k}$ -algebra A with unit. Let $\phi : A \rightarrow \mathbf{k}$ be a nonzero $\mathbf{k}$ -functional on A . Set $\langle \alpha, \beta \rangle_\phi = \phi(\alpha \beta)$ for $\alpha, \beta \in A$. If it is nondegenerate, then $(A, \langle \cdot, \cdot \rangle_\phi)$ is a $\mathbf{k}$ -Fröbenius algebra. Conversely, given any Fröbenius algebra $(A, \langle \cdot, \cdot \rangle)$, let $\phi(\alpha) = \langle \alpha, e \rangle$, for $\alpha \in A$, then $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_\phi$. Now on $\Lambda(V^*)$, consider a Berezin integral $\int : \Lambda(V^*) \rightarrow \mathbf{k}$ (i.e., a $\mathbf{k}$ -linear functional which is only nonzero

on $\Lambda^{top}(V^*)$). Then it is clear that $\langle \alpha, \beta \rangle = \int \alpha \wedge_w \beta$ defines a structure of Fröbenius algebra on $(\Lambda(V^*), \wedge_w)$.

2. Quantum de Rham complex. Let (P, w) be a Poisson manifold, with bivector field w , whose Schouten-Nijenhuis bracket vanishes. (See Vaisman [9] for definitions.) The fiberwise quantum exterior multiplication defines

$$\wedge_h : \Omega_h(M) \otimes \Omega_h(M) \rightarrow \Omega_h(M),$$

$$\wedge_h : \Omega_{h,h^{-1}}(M) \otimes \Omega_{h,h^{-1}}(M) \rightarrow \Omega_{h,h^{-1}}(M),$$

where $\Omega_h(V^*) = \Omega(V^*)[h] = \Omega(V^*) \otimes_{\mathbf{k}} \mathbf{k}[h]$ and $\Omega_{h,h^{-1}}(V^*) = \Omega(V^*)[h, h^{-1}] = \Omega(V^*) \otimes_{\mathbf{k}} \mathbf{k}[h, h^{-1}]$. Koszul [5] defined an operator $\delta : \Omega^k(M) \rightarrow \Omega^{k-1}(M)$ by

$$\delta\alpha = w \lrcorner d\alpha - d(w \lrcorner \alpha),$$

for $\alpha \in \Omega^k(M)$. He also showed that $\delta^2 = 0$, $d\delta + \delta d = 0$. We define the quantum exterior differential $d_h = d - h\delta : \Omega(M)[h] \rightarrow \Omega(M)[h]$, and similarly on $\Omega(M)[h, h^{-1}]$. Then it is easy to see that $d_h^2 = 0$. One of the technical results in Cao-Zhou [3] is the following

Theorem 2. *For any Poisson manifold (M, w) , d_h satisfies*

$$(3) \quad d_h(\alpha \wedge_h \beta) = (d_h\alpha) \wedge_h \beta + (-1)^{|\alpha|} \alpha \wedge_h (d_h\beta),$$

for α, β in $\Omega(M)[h]$, or α, β in $\Omega(M)[h, h^{-1}]$.

This is proved first for $\deg(\alpha) = 1$ by brute force, then by induction on $\deg(\alpha)$. For regular Poisson manifolds (e.g., symplectic manifolds), there is an easier proof. On such Poisson manifolds, there always exists a torsion-less connection ∇ which preserves w . Then for any local frame $\{e^1, \dots, e^n\}$, and $\alpha \in \Omega(M)$, we have

$$d_h\alpha = e^i \wedge_h \nabla_{e_i} \alpha.$$

This is the analogue of a similar expression for $d + d^*$ in Riemannian geometry (Lawson-Michelsohn [6], Lemma II.5.13). Use the analogue of normal coordinates, the proof of Theorem 2 reduces to the associativity of the quantum exterior multiplication.

3. Quantum de Rham cohomology. For any Poisson manifold (M, w) , the (polynomial) quantum deRham cohomology is defined by

$$Q_h H_{dR}^*(M) = \text{Ker } d_h / \text{Im } d_h,$$

for the quantum exterior differential $d_h : \Omega(M)[h] \rightarrow \Omega(M)[h]$. The Laurent quantum de Rham cohomology is

$$Q_{h,h^{-1}} H_{dR}^*(M) = \text{Ker } d_h / \text{Im } d_h,$$

for $d_h : \Omega(M)[h, h^{-1}] \rightarrow \Omega(M)[h, h^{-1}]$. As a consequence of Theorem 1 and Theorem 2, we have

Theorem 3. *The quantum de Rham cohomology $Q_h H_{dR}^*(M)$ of a Poisson manifold has the following properties:*

$$\begin{aligned} \alpha \wedge_h \beta &= (-1)^{|\alpha||\beta|} \beta \wedge_h \alpha, \\ (\alpha \wedge_h \beta) \wedge_h \gamma &= \alpha \wedge_h (\beta \wedge_h \gamma), \end{aligned}$$

for $\alpha, \beta, \gamma \in Q_h H_{dR}^*(M)$. Similar results hold for Laurent quantum deRham cohomology.

The complex $(\Omega(M)[h], d_h)$ can be regarded as a double complex $(C^{p,q}, -h\delta, d)$, where $C^{p,q} = h^p \Omega^{q-p}(M)$, $p \geq 0$. This is the analogue of Brylinski's double complex $\mathcal{C}_\bullet(M)$ ([2], §1.3). By the standard theory for double complex (Bott-Tu [1], §14), there are two spectral sequences E and E' abutting to $H^*(\Omega[h], d_h) = Q_h H_{dR}^*(M)$, with $E_1^{p,q} = h^p H^q(C^{p,*}, d) = h^p H_{dR}^{q-p}(M)$, $(E'_1)^{p,q} = h^p H^*(C^{*,q}, \delta) = h^p PH_{q-p}(M)$, $p \geq 0$. Since nontrivial $E_1^{p,q}$ all have $p+q$ even, and the differential d_r change the parity of $p+q$, it is routine to prove the following

Theorem 4. *For a Poisson manifold with odd Betti numbers all vanishing, the spectral sequence E degenerate at E_1 , i.e. $d_r = 0$ for all $r \geq 0$, hence $Q_h H_{dR}^*(M)$ is a deformation quantization of $H_{dR}^*(M)$.*

Brylinski [2] proved that on closed Kähler manifold (M, ω) , every de Rham cohomology class has a representative α such that $d\alpha = 0$, $\delta\alpha = 0$. This implies that

Theorem 5. *For a closed Kähler manifold M , the spectral sequence E degenerate at E_1 , i.e. $d_r = 0$ for all $r \geq 0$, hence $Q_h H_{dR}^*(M)$ is a deformation quantization of $H_{dR}^*(M)$.*

Similarly, we regard $(\Omega(M)[h, h^{-1}], d - h\delta)$ as a double complex $(\tilde{C}^{p,q}, -h\delta, d)$, where $\tilde{C}^{p,q} = h^p \Omega^{q-p}(M)$, $p, q \in \mathbb{Z}$. This is essentially Brylinski's double complex $\mathcal{C}^{per}$, but with a different bi-grading. We get two spectral sequences $\tilde{E}$ and $\tilde{E}'$ abutting to $Q_{h, h^{-1}} H_{dR}^*(M)$, with $\tilde{E}_1^{p,q} = h^p H_{dR}^{q-p}(M)$, $(\tilde{E}'_1)^{p,q} = h^p H^*(C^{*,q}, \delta) = h^p PH_{q-p}(M)$, $p, q \in \mathbb{Z}$. It is clear that analogue of Theorem 4 holds for $\tilde{E}$. On the other hand, by a method of Brylinski [2], one can prove the following:

Theorem 6. *For any compact symplectic manifold without boundary, the spectral sequences $\tilde{E}$ and $\tilde{E}'$ degenerate at $\tilde{E}_1$ and $\tilde{E}'_1$ respectively. Hence $Q_{h, h^{-1}} H_{dR}^*(M)$ is a Laurent deformation quantization of $H_{dR}^*(M)$.*

Fixing an isomorphism $H_{dR}^{2n}(M) \cong \mathbb{R}$ then defines a structure of Fröbenius algebra on $(H_{dR}^*(M), \wedge_w)$.

4. Quantum Hard Lefschetz Theorem. For a closed symplectic manifold, the analogue of Hard Lefschetz Theorem [4] holds for $Q_{h, h^{-1}} H_{dR}^*(M)$. We begin with a $2n$ -dimensional symplectic vector space (V, ω) . Brylinski [2] defined a symplectic star operator $*$: $\Lambda^k(V^*) \rightarrow \Lambda^{2n-k}(V^*)$. We can extend it to $\Lambda_{h, h^{-1}}$ by setting $*h = h^{-1}$, and $*h^{-1} = h$. Define $L_h : \Lambda_{h, h^{-1}}(V^*) \rightarrow \Lambda_{h, h^{-1}}(V^*)$ by $L_h(\alpha) = \omega \wedge_h \alpha$. Define $L_h^* = - * L *$, and $A_h : \Lambda_{h, h^{-1}}(V^*) \rightarrow \Lambda_{h, h^{-1}}(V^*)$ by $A_h(\alpha) = (n-k)\alpha$, for $\alpha \in \Lambda_{h, h^{-1}}^{[k]}(V^*)$. Then we have

Lemma 1. *The following identities hold:*

$$[L_h, L_h^*] = 0, \quad [L_h, A_h] = 2L_h, \quad [L_h^*, A_h] = -2L_h^*.$$

Furthermore, if we regard multiplications by h and h^{-1} as operators, then we have

$$[h, h^{-1}] = 0, \quad [L_h, h^{\pm 1}] = [L_h^*, h^{\pm 1}] = 0, \quad [A_h, h^{\pm 1}] = \pm 2h^{\pm 1}.$$

Thus we cannot use the representation theory of $sl(2, \mathbb{C})$ as in the classical theory. Notice that multiplication by h is an isomorphism, whose inverse is multiplication by h^{-1} . Let $M_h = h^{-1}L_h$, $M_h^* = hL_h^*$, then M_h, M_h^*, A_h form an abelian algebra. Notice that it now suffices to find the eigenvalues of M_h on $\Lambda_{h, h^{-1}}^{[0]}(V^*)$ and

$\Lambda_{h,h^{-1}}^{[1]}(V^*)$. In Cao-Zhou [3], this is done by induction on the dimension of V and the following

Lemma 2. *Let $\{M_n\}$ be a sequence of square matrices with coefficient in $\mathbf{k}$ obtained in the following way:*

$$M_{n+1} = \begin{pmatrix} M_n & -I \\ I & M_n + 2I \end{pmatrix}$$

for $n \geq 1$. where I is the identity matrix of the same size as M_n . For any $\lambda \in \mathbf{k}$, and $n \geq 1$, we have

$$\det(M_{n+1} + \lambda I) = \det[M_n + (\lambda + 1)I]^2.$$

Therefore, the eigenvalues of M_{n+1} can be obtained by adding 1 to that of M_n , with the multiplicities doubled.

As a consequence, the eigenvalues of M_h on $\Lambda_{h,h^{-1}}(V^*)$ are n and $n \pm \sqrt{5}/2$, when $\dim(V) = 2n$. For M_h^* , they are $-n$ and $-n \pm \sqrt{5}/2$. Therefore, we have

Theorem 7. *For a symplectic vector space V , the operators L_h and L_h^* are isomorphisms. Furthermore, $\Lambda_{h,h^{-1}}(V^*)$ decomposes into one dimensional eigenspaces of $h^{-1}L_h$ (or hL_h^*) with nonzero eigenvalues.*

Lemma 3. *On a symplectic manifold (M, ω) , we have*

$$[L_h, d_h] = 0, \quad [L_h^*, d_h] = 0, \quad [A_h, d_h] = -d_h.$$

As a consequence, we get:

Theorem 8. *(Quantum Hard Lefschetz Theorem) For any symplectic manifold (M^{2n}, ω) , its Laurent quantum deRham cohomology $Q_{h,h^{-1}} H_{dR}^*(M)$ decomposes into one-dimensional eigenspaces of the operator $h^{-1}L_h$ (or hL_h^*) with nonzero eigenvalues. In particular, L_h and L_h^* are isomorphisms.*

5. Complexified quantum exterior algebra. We also consider real vector space V with an almost complex structure $J \in \text{End}(V)$ such that $J^2 = -Id$. There is an induced linear transformation $\Lambda^2 J : \Lambda^2(V) \rightarrow \Lambda^2(V)$. For any bi-vector $w \in \Lambda^2(V)$, J is said to preserve w if $\Lambda^2 J(w) = w$. Given any bi-vector w which is preserved by J , we can define the quantum exterior product on $\Lambda_h(V^*)$ as in the last section. Now if we tensor everything by $\mathbb{C}$, we get a complex algebra $\mathbb{C}\Lambda_h(V^*)$, which is a deformation quantization of $\mathbb{C}\Lambda(V^*) := \Lambda(V^*) \otimes_{\mathbb{R}} \mathbb{C} = \Lambda_{\mathbb{C}}(V^* \otimes_{\mathbb{R}} \mathbb{C})$. As in complex geometry, we can exploit a natural decomposition as follows. J can be uniquely extended to a complex linear endomorphism, denoted also by J , of $\mathbb{C}V$ also satisfying $J^2 = -Id$. There is a natural identification of complex vector spaces $\mathbb{C}V \cong V^{1,0} \oplus V^{0,1}$, where $V^{1,0}$ and $V^{0,1}$ are eigenspaces of J with eigenvalues $\sqrt{-1}$ and $-\sqrt{-1}$ respectively. As a consequence, there are decompositions

$$\begin{aligned} \mathbb{C}\Lambda(V) &= \oplus_{p,q} \Lambda^{p,q}(V), \\ \mathbb{C}\Lambda(V^*) &= \oplus_{p,q} \Lambda^{p,q}(V^*), \end{aligned}$$

where $\Lambda^{p,q}(V) \cong \Lambda_{\mathbb{C}}^p(V^{1,0}) \otimes_{\mathbb{C}} \Lambda_{\mathbb{C}}^q(V^{0,1})$, and $\Lambda^{p,q}(V^*) \cong \Lambda_{\mathbb{C}}^p((V^{1,0})^*) \otimes_{\mathbb{C}} \Lambda_{\mathbb{C}}^q((V^{0,1})^*)$. We give $\mathbb{C}\Lambda_h(V^*)$ the following $\mathbb{Z} \times \mathbb{Z}$ -bigrading: elements in $\Lambda^{p,q}(V^*)$ has bi-degree (p, q) , and h has bi-degree $(1, 1)$. Since w is preserved by J , it belongs to $\Lambda^{1,1}(V)$

after complexification. Denote by $\Lambda_h^{[p,q]}(V^*)$ the space of homogeneous elements of bi-degree (p, q) . It is then straightforward to see that

$$\Lambda_h^{[p,q]}(V^*) \wedge_h \Lambda_h^{[r,s]}(V^*) \subset \Lambda_h^{[p+r, q+s]}(V^*).$$

Now let ω be a symplectic form on V which is compatible with an almost complex structure on V . Namely, rank of ω is $2n = \dim(V)$, $w(J\cdot, J\cdot) = \omega(\cdot, \cdot)$, and $g(\cdot, \cdot) := \omega(\cdot, J\cdot)$ is a positive definite element of $S^2(V^*)$. Then ω induces a natural Hermitian metric H on $\mathbb{C}\Lambda(V^*)$, such that

$$H(\alpha \wedge_w \beta, \gamma) = H(\alpha, \beta \wedge_w \gamma)$$

for any $\alpha, \beta, \gamma \in \mathbb{C}\Lambda(V^*)$. Here $w \in \Lambda^2(V)$ is obtained from ω by “raising the indices” (For details, see Cao-Zhou [3], §1.5). This shows that the algebra $(\mathbb{C}\Lambda(V^*), \wedge_w)$ has a structure of Hermitian Fräbenius algebra.

6. Quantum Dolbeault cohomology.

On a complex manifold (M, J) with a Poisson structure w , such that J preserves w , define $\delta^{-1,0} : \Omega^{p,q}(M) \rightarrow \Omega^{p-1,q}(M)$ and $\delta^{0,-1}(M) : \Omega^{p,q}(M) \rightarrow \Omega^{p,q-1}(M)$ by

$$\begin{aligned} \delta^{0,-1}\alpha &= w \lrcorner (\partial\alpha) - \partial(w \lrcorner \alpha), \\ \delta^{-1,0}\alpha &= w \lrcorner (\bar{\partial}\alpha) - \bar{\partial}(w \lrcorner \alpha), \end{aligned}$$

for $\alpha \in \Omega^{p,q}(M)$. Set $\partial_h = \partial - h\delta^{0,-1}$, and $\bar{\partial}_h = \bar{\partial} - h\delta^{-1,0}$. Then $d_h = \partial_h + \bar{\partial}_h$. Now $0 = d_h^2 = \partial_h^2 + (\partial_h\bar{\partial}_h + \bar{\partial}_h\partial_h) + \bar{\partial}_h^2$, since they have bi-degrees $(2, 0)$, $(1, 1)$ and $(0, 2)$ respectively. Hence, we have

$$\partial_h^2 = 0, \quad \partial_h\bar{\partial}_h + \bar{\partial}_h\partial_h = 0, \quad \bar{\partial}_h^2 = 0.$$

We then define

$$\begin{aligned} Q_h H^{p,*}(M) &= H(\Omega_h^{[p,*]}(M), \bar{\partial}_h), \\ Q_{h,h^{-1}} H^{p,*}(M) &= H(\Omega_{h,h^{-1}}^{[p,*]}(M), \bar{\partial}_h). \end{aligned}$$

They will be called *quantum Dolbeault cohomology* and *Laurent quantum Dolbeault cohomology* respectively. Several relevant spectral sequences and their degeneracy are considered in Cao-Zhou [3].

7. Quantum integral and quantum Stokes Theorem

Let (M, ω) be a closed $2n$ -dimensional symplectic manifold. Define an integral $\int_h : \Omega_h(M) \rightarrow \mathbb{R}[h]$ as follows. For any $\alpha \in \Omega^j(M)$, if j is odd, set $\int_h \alpha = 0$; if $j = 2n - 2k$ for some integer k , set

$$\int_h \alpha = \int_M \alpha \wedge \frac{\omega^k}{k!}.$$

Extend $\int_h$ to $\Omega_h(M)$ as a $\mathbb{R}[h]$ -module map. We call $\int_h$ the *quantum integral*. Then we have

Theorem 0.1. (*Quantum Stokes Theorem*) For any $\alpha \in \Omega^j(M)$, we have $\int_h d\alpha = 0$, $\int_h h\delta\alpha = 0$, and therefore

$$\int_h d_h \alpha = 0.$$

8. Quantum Chern-Weil theory.

Given a real or complex vector bundle $E \rightarrow M$ over a Poisson manifold M , and a connection ∇^E on it, we define the quantum covariant derivative

$$d_h^{\nabla^E} : \Omega_h^*(E) \rightarrow \Omega_h^*(E)$$

as follows. Let $\mathbf{s}$ be a local frame of E and θ be the connection 1-form in this frame. Then we have: $\nabla \mathbf{s} = \mathbf{s} \otimes \theta$, i.e.,

$$\nabla \mathbf{s}_j = \sum_{k=1}^n \mathbf{s}_k \otimes \theta_j^k.$$

For $\alpha = \mathbf{s} \otimes \phi$, where ϕ is some vector valued form, we define

$$d_h^{\nabla^E} \alpha = \mathbf{s} \otimes (\theta \wedge_h \phi + d_h \phi) = \sum \mathbf{s}_k \otimes (\theta_j^k \wedge_h \phi^j + d_h \phi^k).$$

It is straightforward to check that the definition of $d_h^{\nabla^E}$ is independent of the choice of the local frames (Cao-Zhou [3], Lemma 6.1). Furthermore, there is an element $R_h^E \in \Omega_h^2(\text{End}(E))$, such that for each $k \geq 0$, $(d_h^{\nabla^E})^2$ on $\Omega_h^k(M)$ is given by $(d_h^{\nabla^E})^2 \Phi = \Phi \wedge_h R_h^E$, for any $\Phi \in \Omega_h^*(E)$. R_h^E is called the quantum curvature of ∇^E . In a local frame, R_h^E is given by

$$F_h = d_h \theta + \theta \wedge_h \theta,$$

where θ is the connection 1-form matrix in the local frame.

If p is a polynomial on the space of $n \times n$ -matrices, such that $p(G^{-1}AG) = p(A)$, for any $n \times n$ matrix A , and invertible $n \times n$ -matrix G , then $p(F_h)$ for different frames patch up to a well-defined element $p(R^E) \in \Omega^*(M)[h]$. Similar to the ordinary Chern-Weil theory, it is easy to see that $d_h p(R^E) = 0$. So it defines a class in $Q_h H_{dR}^*(M)$. The usual construction of transgression operator carries over to show that this class is independent of the choice of the connection ∇^E . In this way, one can define quantum Chern classes, quantum Euler class, etc. We will call them quantum characteristic classes. It is clear that we can repeat the same story in Laurent case.

9. Quantum equivariant de Rham cohomology. Let (M, w) be a Poisson manifold, which admits an action by a compact connected Lie group G , such that the G -action preserves the Poisson bi-vector field w . Let $\mathfrak{g}$ be the Lie algebra of G , $\{\xi_a\}$ a basis of $\mathfrak{g}$ and $\{\Theta^a\}$ be the dual basis in $S^1(\mathfrak{g}^*)$. Denote by ι_a the contraction by the vector field generated by the one parameter group corresponding to ξ_a , and L_a the Lie derivative by the same vector field. Imitating the Cartan model for equivariant cohomology, we consider the operator $D_{hG} = d_h + \Theta^a \iota_a = d - h\delta + \Theta^a \iota_a$ acting on $(S(\mathfrak{g}^*) \otimes \Omega(M))^G[h]$. It is well-known that $d + \Theta^a \iota_a$ maps $(S(\mathfrak{g}^*) \otimes \Omega(M))^G$ to itself. Since the G -action preserves w , it is easy to check that δ also preserves $(S(\mathfrak{g}^*) \otimes \Omega(M))^G$. Therefore, D_{hG} is an operator from $(S(\mathfrak{g}^*) \otimes \Omega(M))^G[h]$ to itself. Now on $(S(\mathfrak{g}^*) \otimes \Omega(M))^G[h]$, we have

$$\begin{aligned} D_{hG}^2 &= d_h^2 + (\Theta^a \iota_a)^2 + \Theta^a (d \iota_a + \iota_a d) - h \Theta^a (\delta \iota_a + \iota_a \delta) \\ &= -h \Theta^a (\delta \iota_a + \iota_a \delta). \end{aligned}$$

Since $\delta = \iota_w d - d\iota_w$, we have

$$\begin{aligned}\delta\iota_a + \iota_a\delta &= \iota_w d\iota_a - d\iota_w\iota_a + \iota_a\iota_w d - \iota_a d\iota_w \\ &= \iota_w d\iota_a - d\iota_a\iota_w + \iota_w\iota_a d - \iota_a d\iota_w \\ &= \iota_w L_a - L_a\iota_w = -\iota_{L_a w} = 0.\end{aligned}$$

Hence, $D_{hG}^2 = 0$. We call the cohomology

$$Q_h H_G^*(M) := H^*((S(\mathfrak{g}^*) \otimes \Omega(M))^G[h], D_{hG})$$

the quantum equivariant de Rham cohomology. Similar definitions can be made using Laurent deformation. We will study quantum equivariant de Rham cohomology in a forthcoming paper.

REFERENCES

- [1] R. Bott, Raoul; L. W. Tu, **Differential forms in algebraic topology**. Graduate Texts in Mathematics, 82. Springer-Verlag, New York-Berlin, 1982.
- [2] J.-L. Brylinski, *A differential complex for Poisson manifolds*, **J. Differential Geom.** **28** (1988), no. 1, 93–114.
- [3] H.-D. Cao, J. Zhou, *Quantum de Rham cohomology*, preprint, April, 1998.
- [4] P. Griffiths, J. Harris, *Principles of algebraic geometry*. Pure and Applied Mathematics. Wiley-Interscience [John Wiley & Sons], New York, 1978.
- [5] J.-L. Koszul, *Crochet de Schouten-Nijenhuis et cohomologie*, **The mathematical heritage of lie Cartan (Lyon, 1984)**. *Astisque* 1985, Numero Hors Serie, 257–271.
- [6] H. B. Lawson, Jr., M.-L. Michelsohn, **Spin geometry**. Princeton Mathematical Series, 38. Princeton University Press, Princeton, NJ, 1989.
- [7] J. Li, G. Tian, *Comparison of the algebraic and the symplectic Gromov-Witten invariants*, preprint, December 1997, available at alg-geom/9712035.
- [8] G. Tian, *Quantum cohomology and its associativity*, in **Current developments in mathematics**, 1995 (Cambridge, MA), 361–401, Internat. Press, Cambridge, MA, 1994.
- [9] I. Vaisman, **Lectures on the geometry of Poisson manifolds**. Progress in Mathematics, 118. Birkhuser Verlag, Basel, 1994.

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843
E-mail address: cao@math.tamu.edu, zhou@math.tamu.edu